

Spiroscytalin, a New Tetramic Acid and Other Metabolites of Mixed Biogenesis from *Scytalidium cuboideum*

By: Arlene A. Sy-Cordero, [Mario Figueroa](#), [Huzefa A. Raja](#), Maria Elena Meza Aviña, [Mitchell P. Croatt](#), Audrey F. Adcock, David J. Kroll, [Mansukh C. Wani](#), [Cedric J. Pearce](#), and [Nicholas H. Oberlies](#)

Arlene A. Sy-Cordero, Mario Figueroa, Huzefa A. Raja, Maria Elena Meza Aviña, Mitchell P. Croatt, Audrey F. Adcock, David J. Kroll, Mansukh C. Wani, Cedric J. Pearce, and Nicholas H. Oberlies. "Spiroscytalin, a New Tetramic Acid and Other Metabolites of Mixed Biogenesis from *Scytalidium cuboideum*." *Tetrahedron*, 2015, 71(47), 8899-8904. PMID: 26525642; PMCID: PMC4624200; doi: 10.1016/j.tet.2015.09.073

Made available courtesy of Elsevier: <https://doi.org/10.1016/j.tet.2015.09.073>



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](#).

***© 2015 Elsevier Ltd. Reprinted with permission. This version of the document is not the version of record. Figures and/or pictures may be missing from this format of the document. ***

Abstract:

Spiroscytalin (**1**), a new tetramic acid that possesses an uncommon spiro-ring fusion between a polyketide-derived octalin ring system and a 2,4-pyrrolidinedione, along with two known compounds, leporin B (**2**) and purpactin A (**3**), were isolated from a solid phase culture of the fungus *Scytalidium cuboideum* (MSX 68345). The molecular connectivity of **1–3** was determined using NMR spectroscopy and mass spectrometry. The relative configurations of **1** and **2** were determined by NOESY experiments. The absolute configuration of **1** was determined by electronic circular dichroism (ECD) via a combination of experimental measurements and computational calculations. While leporin B was known, it displayed activities that had not been reported previously, including cytotoxicity against three human tumour cell lines and antibacterial activity against *Candida albicans* and *Staphylococcus aureus*.

Keywords: Tetramic acid | Leporin B | Purpactin A | *Scytalidium cuboideum*


Article:

1. Introduction

Ongoing studies on filamentous fungi as a source of anticancer leads^{1, 2} has provided our team with a view of the broad structural diversity that can be uncovered.³ Natural products from mixed biogenetic origin^{4, 5, 6, 7, 8} add to this chemical diversity, as they result from the merging of precursors from two distinct biosynthetic pools (i.e., acetylCoA and amino acids). Tetramic acids⁶ represent a good example, due to their heterocyclic 2,4-pyrrolidinedione ring system, and

this compound class has displayed a variety of biological activities, ranging from HIV-1 integrase inhibitors⁵ to antibacterial and antifungal activities⁴ to cytotoxicity.⁶ These activities, in turn, have spawned further interest in discovering new members of the class^{9, 10} and paved the way for synthetic studies of natural products^{11, 12, 13} and derivatives thereof.^{14, 15, 16} A bioactivity-directed purification study on a solid phase culture of *Scytalidium cuboideum* (MSX 68345) afforded three bioactive compounds, two of which were of mixed biogenetic origin. The new compound, spiroscytalin (**1**), was from the tetramic acid class, while one of the known compounds, leporin B (**2**), was an *N*-hydroxy-2-pyridone alkaloid;^{8, 17} the other known compound was purpactin A (**3**).

Structures



2. Results and discussion

Compound **1** was assigned the molecular formula $C_{22}H_{27}NO_2$ via HRESIMS data (m/z 338.2113 for $[M+H]^+$), establishing an index of hydrogen deficiency (IHD) of 10. The 1H NMR data (Table 1, Fig. S1) showed resonances for a monosubstituted aromatic ring (δ_H 7.22–7.35 for H-17 to H-21) as well as a methylene unit (δ_H 3.35 and 2.51 for H₂-15) connected to it by virtue of HMBC correlations, suggestive of a phenylalanine moiety. Two olefinic protons were also observed (δ_H 5.77 and 5.34 for H-12 and H-13, respectively) as well as five methine protons (δ_H 3.78, 2.28, 1.90, 2.46, and 2.73 for H-3, H-6, H-7, H-11, and H-14, respectively), six more methylene protons (δ_H 1.38, 1.27; 1.49 \times 2; and 1.78, 1.55 for H₂-8, H₂-9, and H₂-10, respectively), and two methyl groups that were each doublets (δ_H 1.13 and 0.89, for CH₃-7 and CH₃-14, respectively). Finally, one NH proton was observed at δ_H 6.41. The ^{13}C NMR data (Table 1) showed resonances for 22 carbons, supportive of the molecular formula, including signals for a ketone carbonyl (δ_C 212.2 for C-4), an amide/ester carbonyl (δ_C 177.2 for C-1),

eight aromatic and olefinic carbons (δ_{C} 132.5, 126.2, 136.6, 129.0 \times 2, 129.3 \times 2, and 127.5 for C-12, C-13, C-16, C-17/C-21, C-18/C-20, and C-19, respectively), one nitrogen bearing carbon (δ_{C} 64.2 for C-3), one sp^3 quaternary carbon (δ_{C} 59.0 for C-5), four methine carbons (δ_{C} 43.8, 30.2, 31.6, 38.0 for C-6, C-7, C-11 and C-14, respectively), four methylene carbons (δ_{C} 28.5, 19.6, 27.6 and 38.5 for C-8, C-9, C-10, and C-15, respectively), and two methyl carbons (δ_{C} 21.4 and 16.2 for CH_3 -7 and CH_3 -14, respectively). COSY correlations from H-6 to H-7 and H-11, from H-11 to H-10 and H-12, from H-7 to CH_3 -7, and from H-14 to CH_3 -14, helped set-up the octalin (octahydronaphthalene) ring system (Fig. 1A). Moreover, COSY correlations from the NH proton to H-3, and COSY correlations, as well as coupling constants, between H₂-15 and H-3 established the tetramic acid moiety. The key HMBC correlations (Fig. 1A) from the NH proton to C-5 and C-15, from H-6 to C-1, C-4, C-3, C-7, CH_3 -7, C-10, and C-11, from H-13 to C-5, C-11, C-14, and CH_3 -14, as well as from CH_3 -14 to C-5, C-13, and C-14, suggested the spiro-ring juncture at C-5 between the octalin ring system and the 2,4-pyrrolidinedione ring system. The latter was a common structural motif in tetramic acids, of which many examples reside in the literature.⁶ Although uncommon, the spiro-ring fusion between a polyketide-derived octalin ring system and a 2,4-pyrrolidinedione ring system has precedent in a few compounds, such as spylidone⁷ and delaminomycins.¹⁸ Compound **1** was ascribed the trivial name spiroscytalin.

Table 1. NMR data for compound **1** (500 MHz for ^1H , 125 MHz for ^{13}C ; δ in ppm; CDCl_3)

Position	δ_{H} , m (J in Hz)	δ_{C}	DEPT
1	—	177.2	C
3	3.78, dd (11.5, 3.0)	64.2	CH
4	—	212.2	C
5	—	59.0	C
6	2.28, br d (6.3)	43.8	CH
7	1.90, m	30.2	CH
8	1.38, br d (13.2)/1.27, m	28.5	CH_2
9	1.49, m/1.49, m	19.6	CH_2
10	1.78, m/1.55, m	27.6	CH_2
11	2.46, m	31.6	CH
12	5.77, dt (10.0, 3.0)	132.5	CH
13	5.34, dt (10.0, 2.0)	126.2	CH
14	2.73, m	38.0	CH
15	3.35, dd (14.0, 3.0)/2.51, dd (14.0, 11.5)	38.5	CH_2
16	—	136.6	C
17	7.22, d (7.5)	129.0	CH
18	7.35, t (7.5)	129.3	CH
19	7.29, d (7.5)	127.5	CH
20	7.35, t (7.5)	129.3	CH
21	7.22, d (7.5)	129.0	CH
7- CH_3	1.13, d (7.5)	21.4	CH_3
14- CH_3	0.89, d (7.5)	16.2	CH_3
NH	6.41, s	—	—




Fig. 1. A) COSY and selected HMBC correlations for compound **1**. B) Key NOESY interactions from the ‘front’ and ‘back’ view (180° turn through the x-axis) observed in the spectrum of **1** as noted using the DFT (B3LYP/DGDZVP) minimum energy model for **1**.

Solving the relative configuration of **1** was somewhat challenging, leading to two possibilities (structures **1** or **4**) that could fit the data. Due to overlapping signals in the ^1H NMR spectrum, many of the J -values were not informative. However, a coupling constant of 6.3 Hz between H-6 and H-11 suggested a *cis*-fused octalin ring system,¹⁹ and this was confirmed by a strong NOESY correlation between these two protons (Fig. 1B). Strong NOESY correlations were also observed between H-6 and H-14, indicating a 1,3-pseudodiaxial relationship. A proton decoupling experiment was performed to confirm the J -value between H-6 and H-7. Irradiation of H-11 resulted in H-6 remaining as a broad doublet ($J=5.2$ Hz), indicating a *gauche* relationship between H-6 and H-7; this finding was supported by a strong NOESY correlation between H-6 and CH₃-7. The configuration at C-7 was enabled by a NOESY correlation between CH₃-7 and H-11, indicative of a 1,3-diaxial relationship. Finally, a NOESY correlation between H-3 and CH₃-14 suggested that they were on the same face of the molecule. At this stage, diastereomers **1** and **4** were both consistent with these NMR data, as were their respective enantiomers. Presumably, the configuration of the 2,4-pyrrolidinedione ring system depended upon whether an *l*- or *d*-phenylalanine was incorporated in the molecule.

Compound **1** exhibited an UV profile similar to spylidone,⁷ and tetramic acids such as equisetin,⁵ phomasetin,⁵ and conioisetin⁴ have been shown to display consistent circular dichroism spectra. Thus, electronic circular dichroism (ECD) calculations were carried out to assign the absolute configuration using density functional theory (DFT).^{20, 21, 22, 23} Given the aforementioned relative configuration considerations, two distinct diastereoisomers were possible, either the 3*R*,5*S* (**1**) or 3*S*,5*R* (**4**). The respective enantiomers of these structures are also possible based on the data obtained, however, only structures **1** and **4** were utilized in the calculations since the ECD spectra would simply be antipodal for their enantiomers. Thus, a Monte Carlo conformational search, utilizing the MMFF94 molecular mechanics force field within a 5 kcal/mol energy window, led to three minimum conformers for each diastereoisomer. Reoptimization of the geometries of all conformers using DFT at the B3LYP/DGDZVP level, as implemented in the Gaussian 09 software, lead to the relative free energies and equilibrium

Boltzmann-weighted populations of the diastereoisomeric series incorporating either 3*R*,5*S*-**1** or 3*S*,5*R*-**4** (Fig. 2 and Tables S1 and S2). For further verification, calculations were also performed for the 3*S*,5*S* and 3*R*,5*R* diastereoisomers (data not shown); these two possibilities were ruled out because a NOESY correlation would not be possible from H-3 and CH₃-14. The three energy minimized conformers for the **1** and **4** series were submitted to ECD calculations by time-dependent density functional theory (TDDFT) at the B3LYP/DGDZVP level, considering the lowest 15 excited states with the CPCM solvent model for MeOH. The resulting ECD spectra for each series were energetically Boltzmann-weighted according to their respective conformational distribution and compared with the experimental ECD spectrum (Fig. 3 and Tables S1 and S2). The experimental ECD spectrum (Fig. 3) displayed a positive Cotton effect (CE) in the region of 210–230 nm and a negative CE in the region of 260–300 nm. These data were in good agreement with the CE calculated for the **1** series. Therefore, the independent TDDFT calculation method supported the absolute configuration of **1** as (3*R*,5*S*,6*R*,7*S*,11*R*,14*S*), suggesting the incorporation of L-phenylalanine, which was consistent with biosynthetic considerations.




Fig. 2. DFT (B3LYP/DGDZVP) geometry optimized conformers of **1** and **4** showing equilibrium population percentages (*P*) based on ΔG (Gibbs free energies relative to the absolute *G* value for the global minimum), assuming Boltzmann statistics at *T*=298.15 K and 1 atm.

Compound **2** was assigned the molecular formula C₂₂H₂₅NO₃ on the basis of HRESIMS data (*m/z* 352.1909 for [M+H]⁺), establishing an IHD of 11. The ¹H and ¹³C NMR data (Fig. S2) of **2** matched that of a known compound leporin B,¹⁷ which is a 2-hydroxy-4-pyridone alkaloid and also of mixed biogenetic origin.⁶ Originally isolated from an unidentified fungal strain, **2** was reported as a transcription activator of hexokinase II gene, a possible target for the treatment of type 2 diabetes.²⁴ We attempted to determine the absolute configuration of **2** via crystallography by making a bromobenzoyl analogue. However, this was unsuccessful due to

paucity of material and the complex reactivity of the hydroxamate relative to a simple alcohol. Thus, relative configuration is shown for **2**.




Fig. 3. Comparison of the experimental ECD spectrum of **1** with the calculated ECD spectra of (3*R*,5*S*)-**1** (**1a–c** series) and (3*S*,5*R*)-**4** (**4a–c** series).

Compound **3** was ascribed a molecular formula of $C_{23}H_{26}O_7$ based on HRESIMS data (m/z 437.1563 for $[M+Na]^+$), indicating an IHD of 11. The 1H (Fig. S3) and ^{13}C NMR and optical rotation data of **3** were in excellent agreement with those reported for purpactin A.²⁵ Purpactin A was first isolated from *Penicillium purpurogenum* and was reported to be an inhibitor of acyl-CoA:cholesterol acyltransferase.²⁵

Compounds **1–3** were tested against three human tumour cells lines (Table 2). Compounds **1** and **3** were moderately active in the human tumour panel (i.e., IC_{50} values between 10 and 20 μM against all cell lines). The biological activity of the crude extract was likely due to leporin B (**2**), which displayed potent, sub μM activity against all three cell lines; cytotoxicity for **2** had not been reported previously. All three compounds were also tested for antimicrobial activity, and **2** showed strong activities against *Candida albicans* and *Staphylococcus aureus* (with MIC values of 6.25 and 25 $\mu g/mL$, respectively). Studies are ongoing to increase the supply of **2**, such that it can undergo more advanced testing.

Table 2. Cytotoxicity against a panel of human tumour cell lines of compounds **1–3**

Compound	IC_{50} values (in μM)		
	MCF7	H460	SF268
1	20.5	17.6	21.9
2	0.2	0.1	0.2
3	11.4	17.3	19.3
Camptothecin	0.07	0.01	0.03

3. Conclusion

A series of compounds of mixed biogenetic origin were isolated and characterized from a fungal culture, including a new tetramic acid analogue (**1**). The biological activities were most promising for the known, yet sparsely studied fungal metabolite, leporin B (**2**). The biosynthetic machinery utilized by the fungus to produce such compounds is intriguing, especially from the

standpoint of future studies to optimize the production of **2** and/or generate analogues via epigenetic manipulation of the growth conditions.

4. Experimental

4.1. General experimental procedures

UV spectra and optical rotations were acquired on a Varian Cary 100 Bio UV–vis spectrophotometer and a Rudolph Research Autopol III polarimeter, respectively. NMR spectra were recorded on JEOL ECA-500, operating at 500 MHz for ^1H and 125 MHz for ^{13}C . HRESIMS was performed on a Thermo LTQ Orbitrap XL mass spectrometer equipped with an electrospray ionization source. HPLC was carried out using a Varian Prostar HPLC system equipped with ProStar 210 pumps and a Prostar 335 photodiode array detector, with data collected and analyzed using Galaxie Chromatography Workstation software (version 1.9.3.2). For preparative HPLC, a YMC ODS-A (5 μm ; 250 \times 20 mm) column was used with a 10 mL/min flow rate; for the semi-preparative HPLC, a YMC ODS-A (5 μm ; 250 \times 10 mm) column was used with a 5 mL/min flow rate; for analytical HPLC, a YMC ODS-A (5 μm ; 150 \times 4.6 mm) column was used with a 1 mL/min flow rate (all from Waters). For analytical HPLC, a MetaTherm HPLC column temperature controller (Varian) maintained the temperature at 30 $^{\circ}\text{C}$.

4.2. Producing organism and fermentation

Mycosynthetix fungal strain MSX 68345 was isolated by Dr. Barry Katz of MYCOsearch in 1993 from leaf litter from a hardwood forest. The entire nuclear ribosomal ITS spacer region (ITS1-5.8S-ITS2)²⁶ and first 600 bp of the nuclear ribosomal Large Subunit (LSU) rRNA gene were sequenced using methods outlined previously.²⁷ Prior to BLAST search, the ITS region was edited using recent guidelines for establishing authenticity and reliability of ITS sequence data.²⁸ A BLAST search with the NCBI GenBank database using the ITS region revealed close identity with isolates of *S. cuboideum* (GenBank AB213451; Identities=599/613 (98%), Gaps=11/613; GenBank AB213444.1; Identities=595/612 (97%). Gaps=13/612). *S. cuboideum* represents a new name for a previously described fungus (*Arthrographis cuboidea*) based on recent phylogenetic analyses using SSU, ITS and RPB2 sequences.^{29, 30} To confirm the identification of MSX 68345 using BLAST search, a Maximum Likelihood (ML) analysis was performed using RAxML.³¹ The combined ITS and LSU sequence of MSX 68345 was subjected to a BLAST search and aligned with taxa with highest affinity from the GenBank BLAST search as well as with other searches implemented through the Fungal Barcoding project (<http://www.fungalbarcoding.org/>). The ML analysis indicated that MSX 68345 was phylogenetically related to a clade consisting of several isolates of *S. cuboideum*, including the type strain (CBS 241.62; AB213451, GQ272628) with strong ML bootstrap support (Fig. S4). Based on results obtained through BLAST search and phylogenetic analysis, MSX 68345 was identified as *S. cuboideum* (Helotiales, Leotiomycetes, Pezizomycotina, Ascomycota). The combined ITS-LSU sequence was deposited in GenBank (Accession No. KR920748).

The culture was stored on a malt extract slant and was transferred periodically. A fresh culture was grown on a similar slant, and a piece was transferred to a medium containing 2% soy peptone, 2% dextrose, and 1% yeast extract (YESD media). Following incubation (7 d) at 22 $^{\circ}\text{C}$

with agitation, the culture was used to inoculate 50 mL of a rice medium, prepared using rice to which was added a vitamin solution and twice the volume of rice with H₂O, in a 250 mL Erlenmeyer flask. This was incubated at 22 °C until the culture showed good growth (approximately 14 d). The scale up culture was grown in a 2.8 L Fernbach flask containing 150 g rice and 300 mL H₂O and was inoculated using a seed culture grown in YESD medium. This was incubated at 22 °C for 14 d.

4.3. Extraction and isolation

To the large scale solid fermentation was added 500 mL of 1:1 MeOH/CHCl₃. The mixture was shaken for 24 h, filtered, and CHCl₃ and H₂O were added to generate a biphasic solution of 4:1:5 (CHCl₃:MeOH:H₂O). The solution was stirred for 2 h, transferred to a separatory funnel, and the organic (lower) layer was removed. This organic extract was evaporated and defatted by partitioning between hexanes and CH₃CN (1:1). The CH₃CN partition (494 mg) was fractionated via flash chromatography using silica gel via a hexanes/CHCl₃/MeOH gradient to afford six fractions. Fraction four (46.5 mg), which eluted with 0–5% MeOH in CHCl₃ over 18 min, was separated further by reversed-phase HPLC (20:80–100:0% CH₃CN–H₂O over 100 min) to obtain eight fractions. Fraction two yielded purpactin A (**3**; 8.0 mg), fraction three yielded the new compound spiroscytalin (**1**; 2.4 mg), and fraction four yielded leporin B (**2**; 6.5 mg). Two more scale-ups of the same extract were processed in the same way outlined above to obtain the same compounds with the following combined yields: Compounds **1** (7.5 mg), **2** (20 mg), and **3** (16 mg).

4.3.1. Spiroscytalin (**1**)

White powder; $[\alpha]_D^{25} -7.7$ (*c* 0.01, MeOH); UV (MeOH) λ_{\max} (log ϵ)=376 (6.28); ¹H, ¹³C NMR, and HMBC data, see Table 1; HRESIMS *m/z* 338.2113 for [M+H]⁺, calcd for C₂₂H₂₈NO₂, 338.2115.

4.3.2. Leporin B (**2**)

White powder; $[\alpha]_D^{25} -67.7$ (*c* 0.04, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.60, s (H-3), 7.41, m (H-18/H-22), 7.37, m (H-19/H-21), 7.29, m (H-20), 5.73, dq (*J*=16, 7; H-15), 5.37, ddq (*J*=15, 8.6, 2; H-14), 4.82, dd (*J*=11, 9; H-13), 2.79, dd (*J*=10, 3.5; H-7), 1.77, m (H-8), 1.60, m (H-9a/H-9b), 1.51, m (H-10a), 1.39, tq (*J*=3.5, 13; H-10b), 1.76, m (H-11a), 1.60, m (H-11b), 1.72, m (H-12), 1.71, dd (*J*=7, 2; H₃-16), 0.86, d (*J*=6; H₃-23); ¹³C NMR (125 MHz, CDCl₃) δ 158.2 (C-1), 129.2 (C-3), 111.2 (C-4), 157.6 (C-5), 113.7 (C-6), 38.1 (C-7), 35.9 (C-8), 35.8 (C-9), 20.9 (C-10), 26.5 (C-11), 35.2 (C-12), 78.2 (C-13), 129.8 (C-14), 131.2 (C-15), 17.9 (C-16), 133.7 (C-17), 129.1 (C-18/C-22), 128.3 (C-19/C-21), 127.5 (C-20), 20.4 (C-23); HRESIMS *m/z* 352.1909 for [M+H]⁺, calcd for C₂₂H₂₆NO₃, 352.1907.

4.3.3. Purpactin A (**3**)

White powder; $[\alpha]_D^{25} -46.1$ (*c* 0.01, CHCl₃); HRESIMS *m/z* 437.1563 for [M+Na]⁺, calcd for C₂₃H₂₆O₇Na, 437.1571.

4.4. Computational methods

Theoretical calculations of TDDFT ECD spectra for compound **1** were performed with the Gaussian'03 program package.³² Geometry optimizations for diastereoisomers **1** and **4** were carried out using the MMFF94 molecular mechanics force field calculations as implemented in the Spartan'08 program. A Monte Carlo search protocol³³ was carried out considering an energy cutoff of 5 kcal/mol, and this generated two sets of conformers, which were made up of three conformers each (**1a–1c** and **4a–4c**; Fig. 2). In each case, the minimum energy structures were filtered and checked for duplicity. No additional minimum energy structures were found. The conformers were optimized by DFT calculations at the B3LYP/DGDZVP level of theory, and thermochemical properties, IR, and vibrational analyses were performed at the same level. The self-consistent reaction field method (SCRF) with conductor-like continuum solvent model (COSMO) was employed to perform the ECD calculation of all conformers of **1**- and **4**-series in MeOH with the same basis set. The calculated excitation energy (in nm) and rotatory strength R in both dipole velocity (R_{vel}) and dipole length (R_{len}) forms were simulated into an ECD curve by using the following Gaussian function:

$$\Delta_{\varepsilon}(E) = \sum_{i=1}^n \varepsilon_i(E) = \sum_{i=1}^n \left(\frac{R_i E_i}{2.29 \times 10^{-39} \sqrt{\pi} \sigma} \exp \left[- \left(\frac{E - E_i}{\sigma} \right)^2 \right] \right)$$

where σ was the width of the band at $1/e$ height, and E_i and R_i were the excitation energies and rotatory strengths for transition i , respectively. $\sigma=0.40$ eV and R_{vel} were used.

4.5. Cytotoxicity assay

The cytotoxicity measurements against MCF-7³⁴ human breast carcinoma (Barbara A. Karmanos Cancer Center, Detroit, MI), NCI-H460³⁵ human large cell lung carcinoma (HTB-177, American Type Culture Collection, Manassas, VA), and SF-268³⁶ human astrocytoma (NCI Developmental Therapeutics Program, Frederick, MD) cell lines were performed as described previously.³⁷

4.6. Antimicrobial assay

The compounds were screened for antimicrobial activity using an agar plate diffusion assay as described previously.³⁸

Acknowledgements

This research was supported by P01 CA125066 from the National Cancer Institute/National Institutes of Health, Bethesda, MD. Mycology technical support was provided by Blaise Darveaux and Maurica Lawrence. Assistance from the staff of the NMR (Dr. Franklin J. Moy) and Mass Spectrometry (Dr. Brandie M. Ehrmann) Facilities at the University of North Carolina at Greensboro is gratefully acknowledged. We thank the Dirección General de Asuntos del Personal Académico (DGAPA-UNAM) by grant IB-200213/RR260213, and the Dirección General de Cómputo y de Tecnologías de Información y Comunicación (DGTIC-UNAM), for providing the resources to carry out computational calculations through the KanBalam system.

References

1. Kinghorn, A. D.; Carache de Blanco, E. J.; Chai, H. B.; Orjala, J.; Farnsworth, N. R.; Soejarto, D. D.; Oberlies, N. H.; Wani, M. C.; Kroll, D. J.; Pearce, C. J.; Swanson, S. M.; Kramer, R. A.; Rose, W. C.; Fairchild, C. R.; Vite, G. D.; Emanuel, S.; Jarjoura, D.; Cope, F. O. *Pure Appl. Chem.* 2009, 81, 1051e1063.
2. Orjala, J.; Oberlies, N. H.; Pearce, C. J.; Swanson, S. M.; Kinghorn, A. D. In *Bioactive Compounds from Natural Sources. Natural Products as Lead Compounds in Drug Discovery*; Tringali, C., Ed.; Taylor & Francis: London, UK, 2012; pp 37e63.
3. El-Elimat, T.; Zhang, X.; Jarjoura, D.; Moy, F. J.; Orjala, J.; Kinghorn, A. D.; Pearce, C. J.; Oberlies, N. H. *ACS Med. Chem. Lett.* 2012, 3, 645e649.
4. Segeth, M. P.; Bonnefoy, A.; Bronstrup, M.; Knauf, M.; Schummer, D.; Toti, L.; Vertesy, L.; Wetzal-Raynal, M. C.; Wink, J.; Seibert, G. *J. Antibiot.* 2003, 56, 114e122.
5. Singh, S. B.; Zink, D. L.; Goetz, M. A.; Dombrowski, A. W.; Polishook, J. D.; Hazuda, D. J. *Tetrahedron Lett.* 1998, 39, 2243e2246.
6. Royles, B. J. L. *Chem. Rev.* 1995, 95, 1981e2001.
7. Koyama, N.; Nagahiro, T.; Yamaguchi, Y.; Ohshiro, T.; Masuma, R.; Tomoda, H.; Omura, S. *J. Antibiot.* 2005, 58, 338e345.
8. Jessen, H. J.; Gademann, K. *Nat. Prod. Rep.* 2010, 27, 1168e1185.
9. Kumar, R.; Subramani, R.; Feussner, K. D.; Aalbersberg, W. *Mar. Drugs* 2012, 10, 200e208.
10. Angawi, R. F.; Bavestrello, G.; Calcinai, B.; Dien, H. A.; Donnarumma, G.; Tufano, M. A.; Paoletti, I.; Grimaldi, E.; Chianese, G.; Fattorusso, E.; Taglialatela-Scafati, O. *Mar. Drugs* 2011, 9, 2809e2817.
11. Sengoku, T.; Nagae, Y.; Ujihara, Y.; Takahashi, M.; Yoda, H. *J. Org. Chem.* 2012, 77, 4391e4401.
12. Kempf, K.; Raja, A.; Sasse, F.; Schobert, R. *J. Org. Chem.* 2013, 78, 2455e2461.
13. Athanasellis, G.; Igglessi-Markopoulou, O.; Markopoulos, J. *Bioinorg. Chem. Appl.* 2010, 2010.
14. Castellucci, N.; Gentilucci, L.; Tomasini, C. *Tetrahedron* 2012, 68, 4506e4512.
15. Jeong, Y. C.; Moloney, M. G. *Beilstein J. Org. Chem.* 2013, 9, 1899e1906.

16. Holloway, C. A.; Matthews, C. J.; Jeong, Y.-C.; Moloney, M. G.; Roberts, C. F.; Yaqoob, M. *Chem. Biol. Drug Des.* 2011, 78, 229e235.
17. Zhang, C.; Jin, L.; Mondie, B.; Mitchell, S. S.; Castelhana, A. L.; Cai, W.; Bergenhem, N. *Bioorg. Med. Chem. Lett.* 2003, 13, 1433e1435.
18. Ueno, M.; Someno, T.; Sawa, R.; Iinuma, H.; Naganawa, H.; Ishizuka, M.; Takeuchi, T. *J. Antibiot.* 1993, 46, 1020e1023.
19. Dodziuk, H.; Jaszunski, M.; Schilf, W. *Magn. Reson. Chem.* 2005, 43, 639e646.
20. Bringmann, G.; Bruhn, T.; Maksimenka, K.; Hemberger, Y. *Eur. J. Org. Chem.* 2009, 2009, 2717e2727.
21. Stephens, P. J.; Pan, J. J.; Devlin, F. J.; Urbanova, M.; Hajicek, J. *J. Org. Chem.* 2007, 72, 2508e2524.
22. Stephens, P. J.; Harada, N. *Chirality* 2010, 22, 229e233.
23. Acuna, U. M.; Figueroa, M.; Kavalier, A.; Jancovski, N.; Basile, M. J.; Kennelly, E. J. *J. Nat. Prod.* 2010, 73, 1775e1779.
24. Vestergaard, H.; Bjorbaek, C.; Hansen, T.; Larsen, F. S.; Granner, D. K.; Pedersen, O. J. *Clin. Invest.* 1995, 96, 2639e2645.
25. Nishida, H.; Tomoda, H.; Cao, J.; Okuda, S.; Omura, S. *J. Antibiot.* 1991, 44, 144e151.
26. Schoch, C. L.; Seifert, K. A.; Huhndorf, S.; Robert, V.; Spouge, J. L.; Levesque, C. A.; Chen, W. *Proc. Natl. Acad. Sci. U.S.A.* 2012, 109, 6241e6246.
27. El-Elmat, T.; Figueroa, M.; Raja, H. A.; Graf, T. N.; Adcock, A. F.; Kroll, D. J.; Day, C. S.; Wani, M. C.; Pearce, C. J.; Oberlies, N. H. *J. Nat. Prod.* 2013, 76, 382e387.
28. Nilsson, R. H.; Tedersoo, L.; Abarenkov, K.; Ryberg, M.; Kristiansson, E.; Hartmann, M.; Schoch, C. L.; Nylander, J. A. A.; Bergsten, J.; Porter, T. M.; Jumpponen, A.; Vaishampayan, P.; Ovaskainen, O.; Hallenberg, N.; Bengtsson-Palme, J.; Eriksson, K. M.; Larsson, K.-H.; Larsson, E.; Kõljalg, U. *Mycologia* 2012, 4, 37e63.
29. Kang, H. J.; Sigler, L.; Lee, J.; Gibas, C. F. C.; Yun, S. H.; Lee, Y. W. *Mycologia* 2010, 102, 1167e1184.
30. Giraldo, A.; Sutton, D. A.; Gene, J.; Fothergill, A. W.; Cano, J.; Guarro, J. *Mycopathologia* 2013, 175, 115e121.
31. Stamatakis, A.; Hoover, P.; Rougemont, J. *Syst. Biol.* 2008, 57, 758e771.

32. Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; Montgomery, J. A., Jr.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K. F. R.; Hasegawa, J.; Ishida, M. N., T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A.; Gaussian, I. Eds.: Pittsburgh, PA, 2003.

33. Chang, G.; Guida, W. C.; Still, W. C. *J. Am. Chem. Soc.* 1989, 111, 4379e4386.

34. Soule, H. D.; Vazquez, J.; Long, A.; Albert, S.; Brennan, M. J. *Natl. Cancer Inst.* 1973, 51, 1409e1416.

35. Carney, D. N.; Gazdar, A. F.; Bunn, P. A., Jr.; Guccion, J. G. *Stem Cells* 1982, 1, 149e164.

36. Rosenblum, M. L.; Gerosa, M. A.; Wilson, C. B.; Barger, G. R.; Pertuiset, B. F.; de Tribolet, N.; Dougherty, D. V. *J. Neurosurg.* 1983, 58, 170e176.

37. Li, C.; Lee, D.; Graf, T. N.; Phifer, S. S.; Nakanishi, Y.; Riswan, S.; Setyowati, F. M.; Saribi, A. M.; Soejarto, D. D.; Farnsworth, N. R.; Falkinham, J. O., III; Kroll, D. J.; Kinghorn, A. D.; Wani, M. C.; Oberlies, N. H. *J. Nat. Prod.* 2009, 72, 1949e1953.

38. Ayers, S.; Ehrmann, B. M.; Adcock, A. F.; Kroll, D. J.; Carcache de Blanco, E. J.; Shen, Q.; Swanson, S. M.; Falkinham, J. O., 3rd; Wani, M. C.; Mitchell, S. M.; Pearce, C. J.; Oberlies, N. H. *J. Pept. Sci.* 2012, 18, 500e510.

Supplementary data

Table S1. DFT (B3LYP/DGDZVP) atomic Cartesian coordinates (Å) for conformers 1

Center number	1a				1b				1c		
	Coordinates (Angstroms)				Coordinates (Angstroms)				Coordinates (Angstroms)		
	X	Y	Z		X	Y	Z		X	Y	Z
1	6.567524	1.350420	0.219636		4.813565	2.620234	-1.352140		-4.444670	1.072015	-2.323266
2	5.835402	0.572019	0.019002		4.681172	1.669064	-0.843203		-4.127263	0.693908	-1.354379
3	3.949299	-1.429677	-0.478098		4.346913	-0.778317	0.476106		-3.319821	-0.300251	1.127900
4	4.612305	0.576096	0.696925		3.610817	1.491832	0.034305		-3.435037	-0.520008	-1.274038
5	6.119900	-0.435303	-0.912755		5.585774	0.621442	-1.067366		-4.418631	1.415348	-0.189795
6	5.173045	-1.437885	-1.157666		5.413639	-0.604092	-0.404322		-4.014753	0.911903	1.053433
7	3.651886	-0.425620	0.459751		3.423163	0.266175	0.709415		-3.014931	-1.031106	-0.033842
8	4.405147	1.358106	1.424921		2.911950	2.306489	0.206560		-3.229708	-1.078640	-2.184881
9	7.071011	-0.440656	-1.439267		6.422130	0.759159	-1.747794		-4.961907	2.355266	-0.249912
10	5.387853	-2.226682	-1.874447		6.117715	-1.415126	-0.571074		-4.242596	1.461067	1.963908
11	3.222234	-2.215587	-0.671757		4.232302	-1.723790	1.001376		-3.014406	-0.685901	2.098000
12	2.317429	-0.399511	1.172947		2.273570	0.076103	1.663202		-2.270653	-2.343480	0.053544
13	1.837478	-1.381459	1.137978		2.644160	-0.346594	2.607630		-2.463530	-2.822559	1.019073
14	2.449125	-0.129109	2.226361		1.785942	1.028017	1.886946		-2.637159	-3.032817	-0.717475
15	1.345778	0.645543	0.570178		1.218327	-0.909202	1.103929		-0.727475	-2.297053	-0.089779
16	1.826518	1.636203	0.608757		1.706853	-1.885034	0.940861		-0.379688	-3.339236	0.002631
17	0.000767	0.672233	1.300464		0.593242	-0.395413	-0.189007		-0.014226	-1.415453	0.937942
18	-0.163818	1.187420	2.463971		1.275989	-0.336800	-1.284105		0.077240	-1.710010	2.182237
19	0.932291	0.347757	-0.787235		0.049597	-1.068686	1.947140		-0.243826	-1.731139	-1.334563
20	1.591214	0.240491	-1.550879		0.102269	-1.479694	2.873399		-0.553269	-2.054058	-2.244851
21	-0.391285	0.443701	-1.038554		-1.146050	-1.053476	1.291796		0.830004	-0.918997	-1.234179
22	-0.917835	0.330454	-2.146971		-2.225699	-1.335250	1.813771		1.455586	-0.435873	-2.179563
23	-1.178949	0.797513	0.248646		-0.934916	-0.703372	-0.200973		1.247498	-0.727413	0.247179
24	-1.720220	2.266518	0.158565		-1.253270	-1.950123	-1.105505		2.569544	-1.523600	0.535492
25	-1.869173	2.566833	1.203686		-0.692056	-1.766724	-2.031495		2.540395	-1.713548	1.616292
26	-2.349027	-0.157289	0.677123		-1.709343	0.545409	-0.767035		1.395199	0.735465	0.791159
27	-2.655790	0.282334	1.633549		-1.312459	0.609882	-1.787644		1.593668	0.556205	1.854367
28	-0.731384	3.279150	-0.448612		-0.739663	-3.289599	-0.544625		2.664710	-2.902760	-0.145103
29	-1.153136	4.287109	-0.377382		-0.914920	-4.085316	-1.276523		3.576677	-3.407933	0.189534
30	-0.534487	3.079578	-1.506732		-1.255695	-3.572281	0.378511		2.716100	-2.822093	-1.235591
31	0.221781	3.287895	0.088121		0.336316	-3.260967	-0.347830		1.823057	-3.548986	0.118654
32	-3.070347	2.348319	-0.523247		-2.716609	-2.045487	-1.480230		3.817198	-0.708673	0.264691
33	-3.359454	3.336428	-0.882182		-3.050518	-3.019032	-1.841318		4.740968	-1.280221	0.170814
34	-3.622793	-0.084269	-0.222462		-3.245207	0.351605	-0.943477		2.649532	1.513148	0.280754
35	-4.450922	-0.378463	0.440736		-3.531802	1.049623	-1.745382		2.867241	2.242167	1.076564
36	-3.908961	1.318070	-0.682438		-3.588104	-1.031926	-1.424691		3.854597	0.624429	0.160749
37	-4.861031	1.478893	-1.189523		-4.622615	-1.200221	-1.726840		4.806406	1.116986	-0.041343
38	-1.932933	-1.625801	0.986743		-1.365432	1.901835	-0.083987		0.106372	1.606538	0.730081
39	-0.896944	-1.597656	1.346194		-0.329702	1.828825	0.270433		-0.747664	0.925282	0.806374
40	-3.663198	-1.075323	-1.409848		-4.111123	0.736380	0.279411		2.446458	2.344336	-1.007626
41	-2.962911	-0.745665	-2.181664		-3.978065	-0.011833	1.064991		2.370244	1.670054	-1.864668
42	-4.667531	-1.040568	-1.851877		-5.166666	0.705946	-0.021347		3.340815	2.961887	-1.163410
43	-3.319484	-2.507923	-0.983295		-3.755019	2.124267	0.827326		1.195748	3.228572	-0.935923
44	-3.285443	-3.159736	-1.864919		-4.337882	2.327877	1.734243		1.046850	3.739399	-1.895267
45	-4.109972	-2.906843	-0.334284		-4.034715	2.898220	0.100389		1.336121	4.017317	-0.185242
46	-1.968997	-2.542548	-0.254162		-2.254500	2.210537	1.140202		-0.040777	2.386078	-0.594504
47	-1.182159	-2.242273	-0.957361		-2.023932	1.506890	1.949658		-0.226568	1.684233	-1.416772
48	-1.728858	-3.567715	0.054978		-2.001045	3.208770	1.519514		-0.930872	3.024586	-0.531042
49	-2.776948	-2.198471	2.142951		-1.393107	3.052468	-1.109971		0.027221	2.544484	1.951367
50	-3.848946	-2.211445	1.918933		-2.375530	3.178735	-1.577605		0.871815	3.239677	2.007404
51	-2.638185	-1.606341	3.054701		-0.667643	2.874988	-1.912250		0.012488	1.968217	2.883620
52	-2.475469	-3.228626	2.365426		-1.133105	4.003496	-0.630378		-0.889729	3.144170	1.916307

Table S2. DFT (B3LYP/DGDZVP) atomic Cartesian coordinates (Å) for conformers of **4**

Center number	4a				4b				4c		
	Coordinates (Angstroms)				Coordinates (Angstroms)				Coordinates (Angstroms)		
	X	Y	Z		X	Y	Z		X	Y	Z
1	6.259734	1.340374	-1.191065		4.360166	2.689455	0.898675		-4.669305	-0.232220	2.000003
2	5.591659	0.596757	-0.762857		4.288069	1.681868	0.495821		-4.196373	-0.303127	1.023160
3	3.864959	-1.314086	0.316540		4.092093	-0.895584	-0.558569		-2.981891	-0.509720	-1.480751
4	4.248412	0.562361	-1.152331		3.180454	1.321309	-0.281615		-3.198054	-1.259166	0.801846
5	6.078007	-0.327533	0.171453		5.301890	0.750783	0.750887		-4.594426	0.552539	-0.011810
6	5.209489	-1.284625	0.709019		5.199668	-0.541634	0.219315		-3.983246	0.444975	-1.266769
7	3.363960	-0.392609	-0.618639		3.064070	0.028621	-0.818066		-2.572628	-1.374667	-0.451834
8	3.885013	1.280235	-1.885454		2.399996	2.054223	-0.474631		-2.912166	-1.928978	1.609905
9	7.122394	-0.304103	0.472817		6.163961	1.029460	1.352045		-5.375305	1.290260	0.156733
10	5.577322	-2.009915	1.431007		5.984857	-1.270639	0.405637		-4.287721	1.100244	-2.079622
11	3.200323	-2.066181	0.737149		4.030012	-1.898965	-0.976326		-2.518785	-0.589684	-2.462148
12	1.901291	-0.401481	-1.017039		1.864798	-0.369712	-1.652367		-1.529991	-2.440196	-0.713543
13	1.801952	-0.220781	-2.093154		1.336027	0.515438	-2.015615		-1.850789	-3.383878	-0.253435
14	-0.266108	0.612255	1.689204		-1.475915	-1.649502	-1.306847		1.125510	-1.228154	1.569561
15	-0.522140	0.591365	2.900394		-2.488116	-2.084997	-1.871560		1.595332	-1.146436	2.711134
16	-0.774872	1.480011	-1.739083		0.530071	-0.652656	1.470459		1.119450	-0.953956	-2.003562
17	-1.278526	0.781671	0.555786		-1.433987	-0.925451	0.038606		1.578829	-0.424235	0.348868
18	-2.001172	2.156857	0.622577		-1.919010	-1.843773	1.196741		3.056971	-0.731071	-0.035445
19	-2.682420	2.151606	1.487161		-3.002965	-1.997177	1.083100		3.709229	-0.321962	0.751228
20	-2.393947	-0.326665	0.586959		-2.347058	0.352413	0.052068		1.503157	1.126372	0.606787
21	-3.041975	-0.024292	1.418551		-3.364334	-0.052745	0.113988		2.376347	1.321411	1.240655
22	-1.082354	3.380047	0.779283		-1.270790	-3.237029	1.256759		3.417014	-2.219809	-0.181143
23	-1.680882	4.296476	0.816101		-1.689732	-3.808024	2.092248		4.484446	-2.324807	-0.402711
24	-0.383181	3.465150	-0.055947		-0.189689	-3.170839	1.400471		2.857152	-2.694064	-0.990406
25	-0.512370	3.321707	1.710857		-1.470807	-3.800077	0.340660		3.220191	-2.761852	0.748452
26	-2.850000	2.242875	-0.614602		-1.705366	-1.066739	2.465763		3.336724	0.038247	-1.298010
27	-3.054352	3.225897	-1.033892		-1.504100	-1.618739	3.381656		4.082250	-0.357896	-1.984825
28	-3.275845	-0.255528	-0.698849		-2.130350	1.175460	1.360995		1.745232	1.920013	-0.712703
29	-4.284856	-0.603188	-0.427989		-3.079133	1.683781	1.592111		2.204244	2.882933	-0.438376
30	-3.428195	1.142662	-1.199895		-1.816803	0.299542	2.530166		2.722813	1.225707	-1.603488
31	-4.023090	1.285995	-2.100122		-1.641894	0.790391	3.485973		2.955724	1.701528	-2.554622
32	-1.939928	-1.779101	0.925590		-2.334620	1.243355	-1.226291		0.284956	1.664995	1.414750
33	-1.068992	-1.713654	1.586495		-2.140519	0.594906	-2.087451		-0.032888	0.880615	2.109570
34	-2.829943	-1.165912	-1.880855		-1.063125	2.305512	1.284507		0.469769	2.287502	-1.521452
35	-1.922604	-0.756102	-2.333286		-0.064187	1.861354	1.260875		0.054269	1.383228	-1.974478
36	-3.613959	-1.141645	-2.647415		-1.129097	2.902413	2.202368		0.760243	2.957062	-2.340608
37	-2.567934	-2.602309	-1.420404		-1.260914	3.190701	0.050296		-0.585969	2.946550	-0.629414
38	-2.216173	-3.199997	-2.269933		-0.463214	3.942044	0.007400		-1.486036	3.155435	-1.220230
39	-3.502076	-3.065835	-1.078332		-2.205840	3.743373	0.129742		-0.220016	3.916064	-0.266468
40	-1.518616	-2.610456	-0.303534		-1.243032	2.333650	-1.220061		-0.933449	2.020947	0.541927
41	-0.582815	-2.210073	-0.710196		-0.258389	1.858913	-1.301801		-1.367588	1.100928	0.139072
42	-1.307363	-3.638270	0.017347		-1.363117	2.963892	-2.110220		-1.701903	2.480874	1.175754
43	-3.038814	-2.497884	1.737620		-3.733962	1.852632	-1.458239		0.722156	2.859574	2.289174
44	-3.990431	-2.556836	1.197726		-4.068246	2.469912	-0.617173		1.124662	3.688812	1.696804
45	-3.228422	-1.975529	2.681923		-4.481018	1.064731	-1.605268		1.497104	2.558221	3.002945
46	-2.733050	-3.522642	1.977868		-3.735684	2.486422	-2.352509		-0.127460	3.248006	2.862572
47	1.457375	-1.377652	-0.804460		2.208363	-0.925818	-2.534427		-1.471681	-2.633946	-1.789651
48	-0.403045	0.771939	-0.698152		0.059609	-0.652059	0.245446		0.674776	-0.962383	-0.770807
49	0.974468	0.494261	1.147413		-0.207937	-1.750573	-1.782438		0.112013	-2.057994	1.195560
50	1.794399	0.440570	1.743656		-0.019049	-2.258429	-2.641110		-0.204695	-2.775753	1.840358
51	1.071701	0.691919	-0.303861		0.864547	-1.279338	-0.897622		-0.074568	-2.195053	-0.254114
52	1.555009	1.660932	-0.507868		1.425961	-2.144163	-0.510691		0.474261	-3.090529	-0.594712




Fig. S1. ^1H (top) and ^{13}C (bottom) NMR spectra of compound **1** [500 MHz for ^1H and 125 MHz for ^{13}C ; CDCl_3].




Fig. S2. ^1H (top) and ^{13}C NMR (bottom) spectra of compound **2** [500 MHz for ^1H and 125 MHz for ^{13}C ; CDCl_3].




Fig. S3. ^1H (top) and DEPTQ (bottom) NMR spectra of compound **3** [500 MHz for ^1H and 100 MHz for ^{13}C ; CDCl_3].




Fig. S4. Phylogram of the most likely tree ($-\ln L = 3472.03$) from a RAxML analysis of 33 taxa based on complete ITS and partial region of the 28S large subunit nrDNA (1007 bp). Numbers refer to RAxML bootstrap support values $\geq 70\%$ based on 1000 replicates. MSX 68345 is highlighted in gray and the *Scytalidium cuboideum* clade with 100% bootstrap value is highlighted yellow. Bar indicates nucleotide substitutions per site.